

## The Extended Mean Values: Definition, Properties, Monotonicities, Comparison, Convexities, Generalizations, and Applications

Feng Qi

Department of Mathematics, Jiaozuo Institute of Technology,  
Jiaozuo City, Henan 454000, China

e-mail address: qifeng@jzit.edu.cn or qifeng618@hotmail.com

URL address: <http://rgmia.vu.edu.au/qi.html>

**ABSTRACT.** Mean values play important roles in the theory of inequalities, and even in the whole of mathematics, since many norms in mathematics are always means. Study of the extended mean values  $E(r, s; x, y)$  is not only interesting but important, both because most the two-variable mean values are special cases of  $E(r, s; x, y)$ , and because it is challenging to study a function whose formulation is so indeterminate.

In this expository article, we summarize the recent main results regarding the study of  $E(r, s; x, y)$ , including its definition, basic properties, monotonicities, comparison, logarithmic convexities, Schur-convexities, generalizations of concepts of mean values, applications to quantum, to theory of special functions, to establishment of Steffensen pairs, and to generalization of Hermite-Hadamard's inequality.

---

2000 *Mathematics Subject Classification.* 05A19, 26A48, 26A51, 26B25, 26D07, 26D10, 26D15, 26D20, 33B20, 41A55, 44A10, 60E15.

*Key words and phrases.* Extended mean values, generalized weighted mean values, generalized abstracted mean values, logarithmic convex, Schur-convex, monotonicity, comparison, definition, recurrence formula, integral expression, gamma function, incomplete gamma function, Steffensen pairs, Hermite-Hadamard inequality, absolutely (completely, regularly) monotonic (convex) function, arithmetic mean of function, quantum, Bernoulli's numbers, Bernoulli's polynomials.

The author was supported in part by NNSF (#10001016) of China, SF for the Prominent Youth of Henan Province, SF of Henan Innovation Talents at Universities, NSF of Henan Province (#004051800), SF for Pure Research of Natural Science of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, China.

The original manuscript is a seminar report giving at the RGMIA on December 10, 2001.

## Contents

<b>1</b>	<b>Definition and Expressions of the Extended Mean Values</b>	<b>64</b>
1.1	Definition of the extended mean values . . . . .	64
1.2	Integral expressions of the extended mean values . . . . .	65
1.3	Inequalities and recurrence formulae for $g(t; x, y)$ . . . . .	66
<b>2</b>	<b>Monotonocities of the Extended Mean Values</b>	<b>67</b>
<b>3</b>	<b>Comparison of the Extended Mean Values</b>	<b>67</b>
<b>4</b>	<b>Convexities of the Extended Mean Values</b>	<b>68</b>
4.1	Definitions of convexities . . . . .	68
4.2	Convexity of the arithmetic mean of a function . . . . .	69
4.3	Logarithmic convexity of the extended mean values . . . . .	70
4.4	Schur-convexity of the extended mean values . . . . .	71
4.4.1	Schur-convexity with $(r, s)$ . . . . .	71
4.4.2	Schur-convexity with $(x, y)$ . . . . .	71
<b>5</b>	<b>Generalizations of Mean Values</b>	<b>72</b>
5.1	Generalized weighted mean values . . . . .	72
5.1.1	Integral case . . . . .	73
5.1.2	Discrete case . . . . .	74
5.2	Generalized abstracted mean values . . . . .	76
5.3	More absolutely monotonic (convex) functions . . . . .	79
<b>6</b>	<b>Applications and Related Results</b>	<b>80</b>
6.1	Application to quantum . . . . .	80
6.2	Generalizations of Bernoulli's numbers and polynomials . . . . .	80
6.3	Generalization of Hermite-Hadamard's inequality . . . . .	80
6.4	Monotonicity results and inequalities involving gamma functions . . . . .	81
6.5	Establishment of Steffensen pairs . . . . .	83

# 1 Definition and Expressions of the Extended Mean Values

The histories of mean values and inequalities are long [9]. The mean values are related to the Mean Value Theorems for the derivative or integral, which are the bridge between the local and global properties of functions. The arithmetic-mean-geometric-mean inequality is probably the most important inequality, and certainly a keystone of the theory of inequalities [2]. Inequalities of mean values are one of the main parts of the theory of inequalities, they have explicit geometric meanings [14]. The theory of mean values plays an important role in the whole mathematics, since many norms in mathematics are often means.

## 1.1 Definition of the extended mean values

In 1975, the extended mean values  $E(r, s; x, y)$  were defined in [51] by K. B. Stolarsky as follows

$$E(r, s; x, y) = \left[ \frac{r}{s} \cdot \frac{y^s - x^s}{y^r - x^r} \right]^{1/(s-r)}, \quad rs(r-s)(x-y) \neq 0; \quad (1.1)$$

$$E(r, 0; x, y) = \left[ \frac{1}{r} \cdot \frac{y^r - x^r}{\ln y - \ln x} \right]^{1/r}, \quad r(x-y) \neq 0; \quad (1.2)$$

$$E(r, r; x, y) = \frac{1}{e^{1/r}} \left[ \frac{x^{r^r} - y^{r^r}}{y^{r^r}} \right]^{1/(x^r - y^r)}, \quad r(x-y) \neq 0; \quad (1.3)$$

$$E(0, 0; x, y) = \sqrt{xy}, \quad x \neq y; \quad (1.4)$$

$$E(r, s; x, x) = x, \quad x = y;$$

where  $x, y > 0$  and  $r, s \in \mathbb{R}$ .

It is easy to see that the extended mean values  $E(r, s; x, y)$  are continuous on the domain  $\{(r, s; x, y) | r, s \in \mathbb{R}; x, y > 0\}$ .

They are symmetric between  $r$  and  $s$  and between  $x$  and  $y$ .

Many basic properties have been researched by E. B. Leach and M. C. Sholander in [19] in 1970's.

Many mean values with two variables are special cases of  $E$ , for example,

$$E(r, 2r; x, y) = M_r(x, y), \quad (\text{power means or Hölder means}) \quad (1.5)$$

$$E(1, p; x, y) = S_p(x, y), \quad (\text{extended logarithmic means}) \quad (1.6)$$

$$E(1, 1; x, y) = I(x, y), \quad (\text{identric or exponential mean}) \quad (1.7)$$

$$E(1, 2; x, y) = A(x, y), \quad (\text{arithmetic mean}) \quad (1.8)$$

$$E(0, 0; x, y) = G(x, y), \quad (\text{geometric mean}) \quad (1.9)$$

$$E(-2, -1; x, y) = H(x, y), \quad (\text{harmonic mean}) \quad (1.10)$$

$$E(0, 1; x, y) = L(x, y). \quad (\text{logarithmic mean}) \quad (1.11)$$

Study of  $E(r, s; x, y)$  is not only interesting but important, both because most of the two-variable mean values are special cases of  $E(r, s; x, y)$ , and because it is challenging to study a function whose formulation is so indeterminate [26].

## 1.2 Integral expressions of the extended mean values

Let

$$g(t) \triangleq g(t; x, y) = \begin{cases} \frac{y^t - x^t}{t}, & t \neq 0; \\ \ln y - \ln x, & t = 0. \end{cases} \quad (1.12)$$

Define a function  $U_n(x; t)$  such that

$$\begin{aligned} U_0(x; t) &= t^x, \\ U_{n+1}(x; t) &= \frac{x \partial U_n(x; t)}{\partial x} - (n+1)U_n(x; t) \end{aligned} \quad (1.13)$$

for  $n$  being a nonnegative integer and  $t > 0$ .

The direct calculation of the  $i$ -th order derivative of  $g(t)$  for  $i \in \mathbb{N}$  is complicated. However, it is easy to see that

$$g^{(i)}(t) = \int_x^y (\ln u)^i u^{t-1} du, \quad y > x > 0, \quad i \in \mathbb{N}. \quad (1.14)$$

Recently, a new expression for the  $i$ -th order derivative of  $g(t; x, y)$  with respect to the variable  $t$  was obtained by the author as follows

$$(-1)^i g^{(i)}(t) = \frac{\Gamma(i+1, -t \ln y) - \Gamma(i+1, -t \ln x)}{t^{i+1}}, \quad (1.15)$$

where  $i$  is a nonnegative integer, and  $\Gamma(z, x)$  denotes the incomplete gamma function defined for  $\text{Re } z > 0$  by

$$\Gamma(z, x) = \int_x^\infty t^{z-1} e^{-t} dt. \quad (1.16)$$

The expressions (1.12), (1.14), and (1.15) of  $g(t; x, y)$  look simple, but they are im-

portant for us. The expression (1.14) can be used to rewrite the extended mean values  $E(r, s; x, y)$  as

$$E(r, s; x, y) = \left( \frac{g(s; x, y)}{g(r; x, y)} \right)^{1/(s-r)}, \quad (r-s)(x-y) \neq 0; \quad (1.17)$$

$$E(r, r; x, y) = \exp \left( \frac{g_r(r; x, y)}{g(r; x, y)} \right), \quad (x-y) \neq 0. \quad (1.18)$$

Taking logarithm in (1.17) and (1.18) yields

$$\ln E(r, s; x, y) = \begin{cases} \frac{1}{s-r} \int_r^s \frac{\partial g(t; x, y)}{\partial t} \cdot \frac{1}{g(t; x, y)} dt, & (r-s)(x-y) \neq 0; \\ \frac{\partial g(r; x, y)}{\partial r} \cdot \frac{1}{g(r; x, y)}, & r = s, x-y \neq 0. \end{cases} \quad (1.19)$$

Note that, the integral expressions (1.14), (1.17), (1.18), and (1.19) of the function  $g$  and the extended mean values  $E(r, s; x, y)$  play key roles in our sequent contents.

### 1.3 Inequalities and recurrence formulae for $g(t; x, y)$

Using Tchebysheff's integral inequality, Hermite-Hadamard's inequality for convex functions and the mathematical induction, some relationships between  $g(x)$  and  $U_n(x, t)$  are deduced, and some recurrence formulae and inequalities of them are given. For examples

**Theorem 1.1** ([46]). *The function  $g(t)$  satisfies*

$$g^{(n)}(t) = \frac{U_n(t; y) - U_n(t; x)}{t^{n+1}}, \quad (1.20)$$

$$\frac{\partial U_n(x, t)}{\partial t} = x^{n+1} (\ln t)^n t^{x-1}. \quad (1.21)$$

**Theorem 1.2** ([46]). *The function  $\frac{g(x+\gamma)}{g(x)}$  is increasing (or decreasing) in  $x$  for  $\gamma > 0$  (or  $\gamma < 0$ ). And  $\left[ \frac{g(x+t)}{g(x)} \right]^{1/t}$ ,  $t \neq 0$ , is increasing with  $t$ .*

**Theorem 1.3** ([46]). *The function  $g(t)$  is absolutely and regularly monotonic on  $\mathbb{R}$  for  $a > 1$ , or on  $(0, \infty)$  for  $b > \frac{1}{a} > 1$ , completely and regularly monotonic on  $\mathbb{R}$  for  $0 < a < b < 1$ , or on  $(-\infty, 0)$  for  $1 < b < \frac{1}{a}$ . Furthermore,  $g(t)$  is absolutely convex on  $\mathbb{R}$ .*

**Theorem 1.4** ([46, 47]). *For  $k, i, j$  being nonnegative integers, we have*

$$g^{(2(i+k)+1)}(t)g^{(2(j+k)+1)}(t) < g^{(2k)}(t)g^{(2(i+j+k-1))}(t). \quad (1.22)$$

The ratio  $\frac{g^{(2(j+k)+1)}(t)}{g^{(2k)}(t)}$  is increasing in  $t$ .

For completeness, we list definitions of absolutely (regularly, completely) monotonic (convex) function as follows.

**Definition 1.1.** A function  $f(t)$  is said to be absolutely monotonic on  $I$  if it has derivatives of all orders and  $f^{(k)}(t) \geq 0$ ,  $t \in I$ ,  $k \in \mathbb{N}$ .

**Definition 1.2.** A function  $f(t)$  is said to be completely monotonic on  $I$  if it has derivatives of all orders and  $(-1)^k f^{(k)}(t) \geq 0$ ,  $t \in I$ ,  $k \in \mathbb{N}$ .

**Definition 1.3.** A function  $f(t)$  is said to be regularly monotonic if it and its derivatives of all orders have constant sign (+ or -; not all the same) on  $I$ .

**Definition 1.4.** A function  $f(t)$  is said to be absolutely convex on  $I$  if it has derivatives of all orders and  $f^{(2k)}(t) \geq 0$ ,  $t \in I$ ,  $k \in \mathbb{N}$ .

The absolutely (completely, regularly) monotonic (convex) functions are useful in Laplace transform [52].

## 2 Monotonocities of the Extended Mean Values

When studying a function, we always consider its monotonicity first. The extended mean values  $E(r, s; x, y)$  are increasing with respect to its all variables. That is

**Theorem 2.1.** *The extended mean values  $E(r, s; x, y)$  is increasing in both  $x$  and  $y$  and in both  $r$  and  $s$ .*

This theorem was verified by E. B. Leach and M. C. Sholander in [20].

Later, using expression (1.17), (1.18) and (1.19), monotonicity of the arithmetic mean of function, Tchebysheff's integral inequality, Cauchy-Schwarz-Buniakowski's inequality and other analytic technique, some simple and new proofs for monotonicity of the extended mean values are provided in [15, 42, 44, 47].

## 3 Comparison of the Extended Mean Values

The comparison of the extended mean values  $E(r, s; x, y)$  is a difficult problem. It was researched in [20]. Five years later, more general results were obtained by Z. Páles in [26] and restated in [25, 29] as follows.

**Theorem 3.1** ([20, 26]). Let  $r, s, u, v$  be real numbers with  $r \neq s$  and  $u \neq v$ , then the inequality

$$E(r, s; a, b) \leq E(u, v; a, b) \quad (3.1)$$

is satisfied for all  $a, b > 0$  if and only if

$$r + s \leq u + v \quad \text{and} \quad e(r, s) \leq e(u, v), \quad (3.2)$$

where

$$e(x, y) = \begin{cases} \frac{x-y}{\ln \frac{x}{y}} & \text{for } xy > 0 \text{ and } x \neq y, \\ 0 & \text{for } xy = 0 \end{cases} \quad (3.3)$$

if either  $0 \leq \min\{r, s, u, v\}$  or  $\max\{r, s, u, v\} \leq 0$ , and

$$e(x, y) = \frac{|x| - |y|}{x - y} \quad \text{for } x, y \in \mathbb{R} \text{ and } x \neq y \quad (3.4)$$

if  $\min\{r, s, u, v\} < 0 < \max\{r, s, u, v\}$ .

## 4 Convexities of the Extended Mean Values

After considering the monotonicity and comparison, it is natural to investigate the convexities of the extended mean values  $E(r, s; x, y)$ .

### 4.1 Definitions of convexities

The concepts of convexities of functions are manifold, for instance, the logarithmically convex and the Schur-convex.

**Definition 4.1** ([24]). A positive function  $f$  defined on an interval  $I$  is logarithmically convex (concave) if its logarithm  $\ln f$  is convex (concave).

**Definition 4.2** ([6, 28]). A function  $f$  with  $n$  arguments on  $I^n$  is Schur-convex on  $I^n$  if  $f(x) \leq f(y)$  for each two  $n$ -tuples  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  in  $I^n$  such that  $x \prec y$  holds, where  $I$  is an interval with nonempty interior.

The relationship of majorization  $x \prec y$  means that

$$\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}, \quad \sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}, \quad (4.1)$$

where  $1 \leq k \leq n-1$  and  $x_{[i]}$  denotes the  $i$ th largest component in  $x$ .

A function  $f$  is Schur-concave if and only if  $-f$  is Schur-convex.

## 4.2 Convexity of the arithmetic mean of a function

The convexities of the (weighted) arithmetic mean of a function (integral arithmetic mean) are important to our proofs for convexities of the extended mean values  $E(r, s; x, y)$ .

The following results can be verified easily.

**Lemma 4.1** ([47]). *If  $f(t)$  is an increasing integrable function on  $I$ , then the arithmetic mean of function  $f(t)$ ,*

$$\phi(r, s) = \begin{cases} \frac{1}{s-r} \int_r^s f(t) dt, & r \neq s, \\ f(r), & r = s, \end{cases} \quad (4.2)$$

is also increasing with both  $r$  and  $s$  on  $I$ .

*If  $f$  is a twice-differentiable convex function, then the function  $\phi(r, s)$  is also convex with both  $r$  and  $s$  on  $I$ .*

In [6], N. Elezović and J. Pečarić proved the following

**Lemma 4.2.** *Let  $f$  be a continuous function on  $I$ . Then the integral arithmetic mean  $\phi(r, s)$  is Schur-convex (Schur-concave) on  $I^2$  if and only if  $f$  is convex (concave) on  $I$ .*

The following necessary and sufficient condition is well-known.

**Lemma 4.3** ([6] and [28, p. 333]). *A continuously differentiable function  $f$  on  $I^2$  (where  $I$  being an open interval) is Schur-convex if and only if it is symmetric and satisfies*

$$\left( \frac{\partial f}{\partial y} - \frac{\partial f}{\partial x} \right) (y - x) > 0 \quad \text{for all } x, y \in I, x \neq y. \quad (4.3)$$

Using Lemma 4.3, we can obtain the Schur-convexities of the weighted arithmetic mean of a function.

**Lemma 4.4** ([45]). *Let  $f$  be a continuous function on  $I$ , let  $p$  be a positive continuous weight on  $I$ . Then the weighted arithmetic mean of a function  $f$  with weight  $p$  defined by*

$$F(x, y) = \begin{cases} \frac{\int_x^y p(t)f(t)dt}{\int_x^y p(t)dt}, & x \neq y, \\ f(x), & x = y \end{cases} \quad (4.4)$$



is Schur-convex (Schur-concave) on  $I^2$  if and only if the inequality

$$\frac{\int_x^y p(t)f(t)dt}{\int_x^y p(t)dt} \leq \frac{p(x)f(x) + p(y)f(y)}{p(x) + p(y)} \quad (4.5)$$

holds (reverses) for all  $x, y \in I$ .

### 4.3 Logarithmic convexity of the extended mean values

By formula (1.19) and Lemma 4.1, we can see that, in order to prove the logarithmic convexity of the extended mean values  $E(r, s; x, y)$ , it suffices to verify the convexity of the function

$$\frac{g'(t)}{g(t)} \triangleq \frac{g'_t(t; x, y)}{g(t; x, y)} \triangleq \frac{\partial g(t; x, y)}{\partial t} \cdot \frac{1}{g(t; x, y)} \quad (4.6)$$

with respect to  $t$ .

Straightforward computation results in

$$\left(\frac{g'(t)}{g(t)}\right)' = \frac{g''(t)g(t) - [g'(t)]^2}{g^2(t)}, \quad (4.7)$$

$$\left(\frac{g'(t)}{g(t)}\right)'' = \frac{g^2(t)g'''(t) - 3g(t)g'(t)g''(t) + 2[g'(t)]^3}{g^3(t)}. \quad (4.8)$$

By a long intricate and standard argument, we obtain the following

**Proposition 4.1** ([33]). *If  $y > x = 1$ , then, for  $t \geq 0$ , we have*

$$g^2(t; 1, y)g_t'''(t; 1, y) - 3g(t; 1, y)g_t'(t; 1, y)g_t''(t; 1, y) + 2[g_t'(t; 1, y)]^3 \leq 0. \quad (4.9)$$

The combination of Proposition 4.1 with equality (4.8) proves that  $\frac{g'_t(t; 1, y)}{g(t; 1, y)}$  is concave on  $[0, \infty)$  with  $t$  for fixed  $y > x = 1$ . Thus, it follows that the extended mean values  $E(r, s; 1, y)$  are logarithmically concave on  $[0, \infty)$  with respect to either  $r$  or  $s$  for  $y > x = 1$ .

By standard arguments, we obtain

$$E(r, s; x, y) = xE\left(r, s; 1, \frac{y}{x}\right), \quad (4.10)$$

$$E(-r, -s; x, y) = \frac{xy}{E(r, s; x, y)}. \quad (4.11)$$

Hence,  $E(r, s; x, y)$  are logarithmically concave on  $[0, \infty)$  with either  $r$  or  $s$  and logarithmically convex on  $(-\infty, 0]$  in either  $r$  or  $s$ , respectively. That is

**Theorem 4.1** ([33]). For all fixed  $x, y > 0$  and  $s \in [0, \infty)$  (or  $r \in [0, \infty)$ , respectively), the extended mean values  $E(r, s; x, y)$  are logarithmically concave in  $r$  (or in  $s$ , respectively) on  $[0, \infty)$ . For all fixed  $x, y > 0$  and  $s \in (-\infty, 0]$  (or  $r \in (-\infty, 0]$ , respectively), the extended mean values  $E(r, s; x, y)$  are logarithmically convex in  $r$  (or in  $s$ , respectively) on  $(-\infty, 0]$ .

#### 4.4 Schur-convexity of the extended mean values

The Schur-convexities are parted into two cases: convexities with respect to  $(r, s)$  and  $(x, y)$ , respectively.

##### 4.4.1 Schur-convexity with $(r, s)$

By the same procedure as the proof of the logarithmic convexity of  $E(r, s; x, y)$  and using Lemma 4.2, we obtain the following

**Theorem 4.2** ([36]). For fixed  $x, y > 0$  and  $x \neq y$ , the extended mean values  $E(r, s; x, y)$  are Schur-concave on  $\mathbb{R}_+^2$  and Schur-convex on  $\mathbb{R}_-^2$  with  $(r, s)$ , where  $\mathbb{R}_+^2$  and  $\mathbb{R}_-^2$  denote  $[0, \infty) \times [0, \infty)$  and  $(-\infty, 0] \times (-\infty, 0]$ , the first and third quadrants, respectively.

Taking  $(r_1, s_1) = (0, 2r)$  and  $(r_2, s_2) = (r, r)$  for  $r \neq 0$ , as a direct consequence of Theorem 4.2, we obtain an inequality between the generalized logarithmic mean values defined by (1.2) and the generalized identity (exponential) mean values defined by (1.3) as follows

**Corollary 4.2.1** ([36]). Let  $x, y > 0$  and  $x \neq y$ . Then, for  $r > 0$ , we have

$$\left[ \frac{1}{2r} \cdot \frac{y^{2r} - x^{2r}}{\ln y - \ln x} \right]^{1/(2r)} \leq \frac{1}{e^{1/r}} \left( \frac{x^{x^r}}{y^{y^r}} \right)^{1/(x^r - y^r)} \quad (4.12)$$

For  $r < 0$ , inequality (4.12) reverses.

##### 4.4.2 Schur-convexity with $(x, y)$

The results on the Schur-convexities with respect to variable  $(x, y)$  are not very perfect. From Lemma 4.4, using the following Theorem 4.4 about inequalities of the arithmetic mean, harmonic mean and logarithmic mean, we have

**Theorem 4.3** ([45]). For fixed point  $(r, s)$  such that  $r, s \notin (0, \frac{3}{2})$  (or  $r, s \in (0, 1]$ , resp.), the extended mean values  $E(r, s; x, y)$  are Schur-concave (or Schur-convex, resp.) with  $(x, y)$  on the domain  $(0, \infty) \times (0, \infty)$ .

As by-products, some inequalities of mean values were established.

**Theorem 4.4 ([45]).** Let  $x, y$  be positive real numbers and  $r \in \mathbb{R}$ .

1. If  $r \leq 0$ , then

$$L(x^r, y^r) \geq [G(x, y)]^r \geq A(x, y)H(x^{r-1}, y^{r-1}). \quad (4.13)$$

the equalities in (4.13) hold only if  $x = y$  or  $r = 0$ .

2. If  $r \geq \frac{3}{2}$ , we have

$$L(x^r, y^r) \geq A(x, y)H(x^{r-1}, y^{r-1}), \quad (4.14)$$

the equality in (4.14) holds only if  $x = y$ .

3. If  $r \in (0, 1]$ , inequality (4.14) reverses without equality unless  $x = y$ .

4. Otherwise, the validity of inequality (4.14) may not be certain.

The results of Theorem 4.4 imply inequalities between the extended mean values and the generalized weighted mean of positive sequence.

**Theorem 4.5 ([45]).** Let  $x, y > 0$ . Then

1. if  $r, s \in (0, 1]$ , we have

$$E(r, s; x, y) \leq M_2((1, 1); (x, y); r - 1, s - 1), \quad (4.15)$$

where  $M_2((1, 1); (x, y); r - 1, s - 1)$  denotes the generalized weighted mean of positive sequence  $(x, y)$  with two parameters  $r - 1$  and  $s - 1$  and constant weight  $(1, 1)$  defined in Definition 5.2;

2. if  $r, s \notin (0, \frac{3}{2})$ , inequality (4.15) reverses;

3. otherwise, the validity of inequality (4.15) may not be certain.

## 5 Generalizations of Mean Values

From (1.14), it is clear that the extended mean values can be rewritten as

$$E(r, s; x, y) = \left( \frac{\int_x^y t^{s-1} dt}{\int_x^y t^{r-1} dt} \right)^{1/(s-r)}. \quad (5.1)$$

## 5.1 Generalized weighted mean values

One of generalizations of mean values, the generalized weighted mean values  $M_{p,f}(r, s; x, y)$ , are classified into two cases.

### 5.1.1 Integral case

It is natural to generalize the concept of the extended mean values  $E(r, s; x, y)$  through replacing  $t$  by a positive function  $f(t)$  and considering a weight in the integrands in (5.1).

**Definition 5.1** ([32, 35]). Let  $x, y, r, s \in \mathbb{R}$ , and  $p(u) \neq 0$  be a nonnegative and integrable function,  $f(u)$  a positive and integrable function on the interval between  $x$  and  $y$ . The generalized weighted mean values, with weight  $p(u)$  and two parameters  $r$  and  $s$ , is defined by

$$M_{p,f}(r, s; x, y) = \left( \frac{\int_x^y p(u) f^s(u) du}{\int_x^y p(u) f^r(u) du} \right)^{1/(s-r)}, \quad (r-s)(x-y) \neq 0; \quad (5.2)$$

$$M_{p,f}(r, r; x, y) = \exp \left( \frac{\int_x^y p(u) f^r(u) \ln f(u) du}{\int_x^y p(u) f^r(u) du} \right), \quad r(x-y) \neq 0; \quad (5.3)$$

$$M_{p,f}(r, 0; x, y) = \left( \frac{\int_x^y p(u) f^r(u) du}{\int_x^y p(u) du} \right)^{1/r}, \quad r(x-y) \neq 0; \quad (5.4)$$

$$M_{p,f}(0, 0; x, y) = \exp \left( \frac{\int_x^y p(u) \ln f(u) du}{\int_x^y p(u) du} \right), \quad x-y \neq 0; \quad (5.5)$$

$$M_{p,f}(r, s; x, x) = f(x).$$

The following lemma is called the revised Cauchy mean value theorem in integral form.

**Lemma 5.1** ([32, 35, 47]). Suppose that  $f(t)$  and  $g(t) \geq 0$  are integrable on  $[a, b]$  and the ratio  $\frac{f(t)}{g(t)}$  has finitely many removable discontinuity points. Then there exists at least one point  $\theta \in (a, b)$  such that

$$\frac{\int_a^b f(t) dt}{\int_a^b g(t) dt} = \lim_{t \rightarrow \theta} \frac{f(t)}{g(t)}. \quad (5.6)$$

Using Lemma 5.1, some basic properties of the generalized weighted mean values  $M_{p,f}(r, s; x, y)$  were yielded as follows.

**Theorem 5.1** ([32]).  $M_{p,f}(r, s; x, y)$  have the following properties

$$m \leq M_{p,f}(r, s; x, y) \leq M, \quad (5.7)$$

$$M_{p,f}(r, s; x, y) = M_{p,f}(r, s; y, x) = M_{p,f}(s, r; x, y), \quad (5.8)$$

$$M_{p,f}^{s-r}(r, s) = M_{p,f}^{s-t}(t, s)M_{p,f}^{t-r}(r, t), \quad (5.9)$$

where  $m = \inf f(u)$ ,  $M = \sup f(u)$ .

In [32] and [44], the monotonicity with  $x$  and  $y$  of  $M_{p,f}(r, s; x, y)$  was proved by three approaches.

**Theorem 5.2.** Let  $p(u) \neq 0$  be a nonnegative and continuous function,  $f(u)$  a positive, increasing (or decreasing, respectively) and continuous function. Then  $M_{p,f}(r, s; x, y)$  increases (or decreases, respectively) with respect to either  $x$  or  $y$ .

Using Cauchy-Schwarz-Buniakowski's inequality, we proved monotonicity of the generalized weighted mean values  $M_{p,f}(r, s; x, y)$  with  $(r, s)$  as follows.

**Theorem 5.3** ([48]). The generalized weighted mean values  $M_{p,f}(r, s; x, y)$  are increasing with both  $r$  and  $s$  for any continuous nonnegative weight  $p$  and continuous positive function  $f$ .

Using Tchebysheff's integral inequality, we have the following two theorems.

**Theorem 5.4** ([32]). Let  $p_1(u) \neq 0$  and  $p_2(u) \neq 0$  be nonnegative and integrable functions on the interval between  $x$  and  $y$ ,  $f(u)$  a positive and integrable function, the ratio  $\frac{p_1(u)}{p_2(u)}$  an integrable function,  $\frac{p_1(u)}{p_2(u)}$  and  $f(u)$  both increasing or both decreasing. Then

$$M_{p_1,f}(r, s; x, y) \geq M_{p_2,f}(r, s; x, y) \quad (5.10)$$

If one of the functions of  $f(u)$  or  $\frac{p_1(u)}{p_2(u)}$  is nonincreasing and the other nondecreasing, then inequality (5.10) is reversed.

**Theorem 5.5** ([32]). Let  $p(u) \neq 0$  be a nonnegative and integrable function, and  $f_1(u)$  and  $f_2(u)$  positive and integrable functions on the interval between  $x$  and  $y$ . If the ratio  $\frac{f_1(u)}{f_2(u)}$  and  $f_2(u)$  are integrable and both increasing or both decreasing, then

$$M_{p,f_1}(r, s; x, y) \geq M_{p,f_2}(r, s; x, y) \quad (5.11)$$

holds for  $r, s \geq 0$  or  $r \geq 0 \geq s$ , and  $\frac{f_1(u)}{f_2(u)} \geq 1$ . The inequality (5.11) is reversed for  $r, s \leq 0$  or  $s \geq 0 \geq r$ , and  $\frac{f_1(u)}{f_2(u)} \leq 1$ .

If one of the functions of  $f_2(u)$  or  $\frac{f_1(u)}{f_2(u)}$  is nonincreasing and the other nondecreasing, then inequality (5.11) is valid for  $r, s \geq 0$  or  $s \geq 0 \geq r$ , and  $\frac{f_1(u)}{f_2(u)} \geq 1$ ; the inequality (5.11) reverses for  $r, s \geq 0$  or  $r \geq 0 \geq s$ , and  $\frac{f_1(u)}{f_2(u)} \leq 1$ .

### 5.1.2 Discrete case

The discrete analogue of the generalized weighted mean values, the generalized weighted mean of positive sequence  $a = (a_1, \dots, a_n)$ , was defined in [31] by

**Definition 5.2.** For a positive sequence  $a = (a_1, \dots, a_n)$  with  $a_i > 0$  and a positive weight  $p = (p_1, \dots, p_n)$  with  $p_i > 0$  for  $1 \leq i \leq n$ , the generalized weighted mean of positive sequence  $a$  with two parameters  $r$  and  $s$  is defined as

$$M_n(p; a; r, s) = \begin{cases} \left( \frac{\sum_{i=1}^n p_i a_i^r}{\sum_{i=1}^n p_i a_i^s} \right)^{1/(r-s)}, & r - s \neq 0; \\ \exp \left( \frac{\sum_{i=1}^n p_i a_i^r \ln a_i}{\sum_{i=1}^n p_i a_i^r} \right), & r - s = 0. \end{cases} \quad (5.12)$$

*Remark 5.1.* For  $s = 0$  we obtain the weighted mean  $M_n^{[r]}(a; p)$  of order  $r$  (see [24]); for  $s = 0, r = -1$ , the weighted harmonic mean; for  $s = 0, r = 0$ , the weighted geometric mean; and for  $s = 0, r = 1$ , the weighted arithmetic mean.

The mean  $M_n(p; a; r, s)$  has some basic properties similar to those of  $M_{p,f}(r, s; x, y)$ , for instance

**Theorem 5.6 ([31]).** The mean  $M_n(p; a; r, s)$  is a continuous function with respect to  $(r, s) \in \mathbb{R}^2$  and has the following properties

$$\begin{aligned} m &\leq M_n(p; a; r, s) \leq M, \\ M_n(p; a; r, s) &= M_n(p; a; s, r), \\ M_n^{s-r}(p; a; r, s) &= M_n^{s-t}(p; a; t, s) \cdot M_n^{t-r}(p; a; r, t), \end{aligned} \quad (5.13)$$

where  $m = \min_{1 \leq i \leq n} \{a_i\}$ ,  $M = \max_{1 \leq i \leq n} \{a_i\}$ .

The inequality property in (5.13) follows from the following elementary inequalities in [24, p. 204] which are due to Cauchy.

For an arbitrary sequence  $b = (b_1, \dots, b_n)$  and a positive sequence  $c = (c_1, \dots, c_n)$ , we have

$$\min_{1 \leq i \leq n} \left\{ \frac{b_i}{c_i} \right\} \leq \frac{\sum_{i=1}^n b_i}{\sum_{i=1}^n c_i} \leq \max_{1 \leq i \leq n} \left\{ \frac{b_i}{c_i} \right\}. \quad (5.14)$$

Equality holds in both of the above inequalities if and only if the sequences  $b$  and  $c$  are proportional.

Using Lemma 4.1 and by standard arguments, we obtain the monotonicity of  $M_n(p; a; r, s)$  with respect to variables  $r$  and  $s$ .

**Theorem 5.7 ([31]).** *The mean  $M_n(p; a; r, s)$  of numbers  $a = (a_1, \dots, a_n)$  with weights  $p = (p_1, \dots, p_n)$  and two parameters  $r$  and  $s$  is increasing in both  $r$  and  $s$ .*

By mathematical induction and inequalities in (5.14), we obtain an inequality for different natural indices  $n$  of  $M_n(p; a; r, s)$ .

**Theorem 5.8 ([31]).** *For a monotonic sequence of positive numbers  $0 < a_1 \leq a_2 \leq \dots$  and positive weights  $p = (p_1, p_2, \dots)$ , if  $m < n$ , then*

$$M_m(p; a; r, s) \leq M_n(p; a; r, s). \quad (5.15)$$

Equality holds if  $a_1 = a_2 = \dots$ .

Using the discrete Tchebysheff's inequality, the following are obtained.

**Theorem 5.9 ([31]).** *Let  $p = (p_1, \dots, p_n)$  and  $q = (q_1, \dots, q_n)$  be positive weights,  $a = (a_1, \dots, a_n)$  a sequence of positive numbers. If the sequences  $(\frac{p_1}{q_1}, \dots, \frac{p_n}{q_n})$  and  $a$  are both nonincreasing or both nondecreasing, then*

$$M_n(p; a; r, s) \geq M_n(q; a; r, s). \quad (5.16)$$

If one of the sequences of  $(\frac{p_1}{q_1}, \dots, \frac{p_n}{q_n})$  or  $a$  is nonincreasing and the other nondecreasing, the inequality (5.16) is reversed.

**Theorem 5.10 ([31]).** *Let  $p = (p_1, \dots, p_n)$  be positive weights,  $a = (a_1, \dots, a_n)$  and  $b = (b_1, \dots, b_n)$  two sequences of positive numbers. If the sequences  $(\frac{a_1}{b_1}, \dots, \frac{a_n}{b_n})$  and  $b$  are both increasing or both decreasing, then*

$$M_n(p; a; r, s) \geq M_n(p; b; r, s) \quad (5.17)$$

holds for  $\frac{a_i}{b_i} \geq 1$ ,  $n \geq i \geq 1$ , and  $r, s \geq 0$  or  $r \geq 0 \geq s$ . The inequality (5.17) is reversed for  $\frac{a_i}{b_i} \leq 1$ ,  $n \geq i \geq 1$ , and  $r, s \leq 0$  or  $s \geq 0 \geq r$ .

If one of the sequences of  $(\frac{a_1}{b_1}, \dots, \frac{a_n}{b_n})$  or  $b$  is nonincreasing and the other nondecreasing, then inequality (5.17) is valid for  $\frac{a_i}{b_i} \geq 1$ ,  $n \geq i \geq 1$  and  $r, s \geq 0$  or  $s \geq 0 \geq r$ ; the inequality (5.17) reverses for  $\frac{a_i}{b_i} \leq 1$ ,  $n \geq i \geq 1$ , and  $r, s \geq 0$  or  $r \geq 0 \geq s$ .

## 5.2 Generalized abstracted mean values

The following definition is an integral analogue of the Definition 3 in [24, p. 75].

**Definition 5.3.** Let  $p$  be a defined, positive and integrable function on  $[x, y]$  for  $x, y \in \mathbb{R}$ ,  $f$  a real-valued and monotonic function on  $[\alpha, \beta]$ . If  $g$  is a function valued on  $[\alpha, \beta]$  and  $f \circ g$  integrable on  $[x, y]$ , the quasi-arithmetic non-symmetrical mean of function  $g$  is defined by

$$M_f(g; p; x, y) = f^{-1} \left( \frac{\int_x^y p(t) f(g(t)) dt}{\int_x^y p(t) dt} \right), \quad (5.18)$$

where  $f^{-1}$  is the inverse function of  $f$ .

*Remark 5.2.* For  $g(t) = t$ ,  $f(t) = t^{r-1}$ ,  $p(t) = 1$ , the mean  $M_f(g; p; x, y)$  reduces to the extended logarithmic means  $S_r(x, y)$ ; for  $p(t) = t^{r-1}$ ,  $g(t) = f(t) = t$ , to the one-parameter mean  $J_r(x, y)$ ; for  $p(t) = f'(t)$ ,  $g(t) = t$ , to the abstracted mean  $M_f(x, y)$ ; for  $g(t) = t$ ,  $p(t) = t^{r-1}$ ,  $f(t) = t^{s-r}$ , to the extended mean values  $E(r, s; x, y)$ ; for  $f(t) = t^r$ , to the weighted mean of order  $r$  of the function  $g$  with weight  $p$  on  $[x, y]$ . If we replace  $p(t)$  by  $p(t)f^r(t)$ ,  $f(t)$  by  $t^{s-r}$ ,  $g(t)$  by  $f(t)$  in (5.18), then we get the generalized weighted mean values  $M_{p,f}(r, s; x, y)$ . Hence, from  $M_f(g; p; x, y)$  we can deduce a lot of the two variable means.

The following properties follow easily from Lemma 5.1 and standard arguments.

**Theorem 5.11 ([31]).** *The mean  $M_f(g; p; x, y)$  has the following properties*

$$\begin{aligned} \alpha &\leq M_f(g; p; x, y) \leq \beta, \\ M_f(g; p; x, y) &= M_f(g; p; y, x), \end{aligned} \quad (5.19)$$

where  $\alpha = \inf_{t \in [x, y]} g(t)$  and  $\beta = \sup_{t \in [x, y]} g(t)$ .

The function  $\frac{1}{x}$  is the reciprocal function of  $f(x) = x$ . Further, we have

**Lemma 5.2 ([31]).** *Suppose the ratio  $\frac{f_1}{f_2}$  is monotonic on a given interval. Then*

$$\left( \frac{f_1}{f_2} \right)^{-1}(x) = \left( \frac{f_2}{f_1} \right)^{-1} \left( \frac{1}{x} \right), \quad (5.20)$$

where  $\left( \frac{f_1}{f_2} \right)^{-1}$  is the inverse function of  $\frac{f_1}{f_2}$ .

These hints remind us that, if replacing  $\frac{1}{s-r}$  by  $\left( \frac{f_1}{f_2} \right)^{-1}$  in Definition 5.2, then we can obtain



**Definition 5.4** ([31]). Let  $f_1$  and  $f_2$  be real-valued functions such that the ratio  $\frac{f_1}{f_2}$  is monotone on the closed interval  $[\alpha, \beta]$ . If  $a = (a_1, \dots, a_n)$  is a sequence of real numbers from  $[\alpha, \beta]$  and  $p = (p_1, \dots, p_n)$  a sequence of positive numbers, the generalized abstracted mean values of numbers  $a$  with respect to functions  $f_1$  and  $f_2$ , with weights  $p$ , is defined by

$$M_n(p; a; f_1, f_2) = \left(\frac{f_1}{f_2}\right)^{-1} \left(\frac{\sum_{i=1}^n p_i f_1(a_i)}{\sum_{i=1}^n p_i f_2(a_i)}\right), \quad (5.21)$$

where  $\left(\frac{f_1}{f_2}\right)^{-1}$  is the inverse function of  $\frac{f_1}{f_2}$ .

The integral analogue of Definition 5.4 is given by

**Definition 5.5** ([31]). Let  $p$  be a positive integrable function defined on  $[x, y]$ ,  $x, y \in \mathbb{R}$ ,  $f_1$  and  $f_2$  real-valued functions and the ratio  $\frac{f_1}{f_2}$  monotone on the interval  $[\alpha, \beta]$ . In addition, let  $g$  be defined on  $[x, y]$  and valued on  $[\alpha, \beta]$ , and  $f_i \circ g$  integrable on  $[x, y]$  for  $i = 1, 2$ . The generalized abstracted mean values of function  $g$  with respect to functions  $f_1$  and  $f_2$  and with weight  $p$  is defined as

$$M(p; g; f_1, f_2; x, y) = \left(\frac{f_1}{f_2}\right)^{-1} \left(\frac{\int_x^y p(t) f_1(g(t)) dt}{\int_x^y p(t) f_2(g(t)) dt}\right), \quad (5.22)$$

where  $\left(\frac{f_1}{f_2}\right)^{-1}$  is the inverse function of  $\frac{f_1}{f_2}$ .

*Remark 5.3.* Set  $f_2 \equiv 1$  in Definition 5.5, then we can obtain Definition 5.3 easily. Replacing  $f$  by  $\frac{f_1}{f_2}$ ,  $p(t)$  by  $p(t)f_2(g(t))$  in Definition 5.3, we arrive at Definition 5.5 directly. Definition 5.3 and Definition 5.5 are equivalent to each other. Analogously, formula (5.21) is equivalent to  $M_f(a; p)$ . Similarly, so are Definition 5.4 and the quasi-arithmetic non-symmetrical mean  $M_f(a; p)$  of numbers  $a = (a_1, \dots, a_n)$  with weights  $p = (p_1, \dots, p_n)$ .

From inequality (5.14), Lemma 5.1, Lemma 5.2 and standard arguments, we have

**Theorem 5.12** ([31]). *The means  $M_n(p; a; f_1, f_2)$  and  $M(p; g; f_1, f_2; x, y)$  have the following properties*

1. Under the conditions of Definition 5.4, we have

$$\begin{aligned} m &\leq M_n(p; a; f_1, f_2) \leq M, \\ M_n(p; a; f_1, f_2) &= M_n(p; a; f_2, f_1), \end{aligned} \quad (5.23)$$

where  $m = \min_{1 \leq i \leq n} \{a_i\}$ ,  $M = \max_{1 \leq i \leq n} \{a_i\}$ ;

2. Under the conditions of Definition 5.5, we have

$$\begin{aligned} \alpha &\leq M(p; g; f_1, f_2; x, y) \leq \beta, \\ M(p; g; f_1, f_2; x, y) &= M(p; g; f_1, f_2; y, x), \\ M(p; g; f_1, f_2; x, y) &= M(p; g; f_2, f_1; x, y), \end{aligned} \quad (5.24)$$

where  $\alpha = \inf_{t \in [x, y]} g(t)$  and  $\beta = \sup_{t \in [x, y]} g(t)$ .

By Lemma 5.1 and standard argument, it follows that

**Theorem 5.13** ([31]). Suppose  $p$  and  $g$  are defined on  $\mathbb{R}$ . If  $f_1 \circ g$  has constant sign and if  $\left(\frac{f_1}{f_2}\right) \circ g$  is increasing (or decreasing, respectively), then  $M(p; g; f_1, f_2; x, y)$  have the inverse (or same) monotonicities as  $\frac{f_1}{f_2}$  with both  $x$  and  $y$ .

The Tchebysheff's integral inequality produces the following two theorems.

**Theorem 5.14** ([31]). Suppose  $f_2 \circ g$  has constant sign on  $[x, y]$ . When  $g(t)$  increases on  $[x, y]$ , if  $\frac{p_1}{p_2}$  is increasing, we have

$$M(p_1; g; f_1, f_2; x, y) \geq M(p_2; g; f_1, f_2; x, y); \quad (5.25)$$

if  $\frac{p_1}{p_2}$  is decreasing, inequality (5.25) reverses.

When  $g(t)$  decreases on  $[x, y]$ , if  $\frac{p_1}{p_2}$  is increasing, then inequality (5.25) is reversed; if  $\frac{p_1}{p_2}$  is decreasing, inequality (5.25) holds.

**Theorem 5.15** ([31]). Suppose  $f_2 \circ g_2$  does not change its sign on  $[x, y]$ .

1. When  $f_2 \circ \left(\frac{g_1}{g_2}\right)$  and  $\left(\frac{f_1}{f_2}\right) \circ g_2$  are both increasing or both decreasing, inequality

$$M(p; g_1; f_1, f_2; x, y) \geq M(p; g_2; f_1, f_2; x, y) \quad (5.26)$$

holds for  $\frac{f_1}{f_2}$  being increasing, or reverses for  $\frac{f_1}{f_2}$  being decreasing.

2. When one of the functions  $f_2 \circ \left(\frac{g_1}{g_2}\right)$  or  $\left(\frac{f_1}{f_2}\right) \circ g_2$  is decreasing and the other increasing, inequality (5.26) holds for  $\frac{f_1}{f_2}$  being decreasing, or reverses for  $\frac{f_1}{f_2}$  being increasing.

### 5.3 More absolutely monotonic (convex) functions

In [31] and [32], some more general absolutely (regularly, completely) monotonic (convex) functions were established, which generalize the related results in [46] restated in Theorem 1.3 of Section 1.3.

**Theorem 5.16** ([32]). Suppose that  $f(u)$  is positive and has derivatives of all orders on the interval  $[a, b]$ . Define  $\psi(t)$  by

$$\psi(t) = \begin{cases} \frac{f'(b) - f'(a)}{t}, & t \neq 0; \\ \ln f(b) - \ln f(a), & t = 0. \end{cases} \quad (5.27)$$

Then

$$\psi^{(n)}(t) = \frac{U_n(t, f(b)) - U_n(t, f(a))}{t^{n+1}}, \quad (5.28)$$

$$\frac{\partial U_n(t, s)}{\partial s} = t^{n+1}(\ln s)^n s^{t-1}, \quad (5.29)$$

where  $U_n$  is defined in (1.13).

**Theorem 5.17** ([32]). If  $f(u) \geq 1$  and  $f'(u) \geq 0$ , then the function  $\psi(t)$  defined by (5.27) is absolutely and regularly monotonic on the interval  $\mathbb{R}$ . If  $0 < f(u) \leq 1$  and  $f'(u) \geq 0$  then  $\psi(t)$  is completely and regularly monotonic on  $\mathbb{R}$ . Moreover,  $\psi(t)$  is absolutely convex on  $\mathbb{R}$ .

**Theorem 5.18** ([31]). Suppose  $F(t) = \int_a^b p(u)f'(u)du$ , where  $t \in \mathbb{R}$ ,  $p(u) \neq 0$  is a non-negative and continuous function, and  $f(u)$  is a positive and continuous function on a given interval  $[a, b]$ . Then

$$F^{(n)}(t) = \int_a^b p(u)f'(u)[\ln f(u)]^n du. \quad (5.30)$$

If  $f(u) \geq 1$ , then  $F(t)$  is absolutely monotone on  $\mathbb{R}$ ; if  $0 < f(u) < 1$ , then  $F(t)$  is completely monotone on  $\mathbb{R}$ . Moreover,  $F(t)$  is absolutely convex on  $\mathbb{R}$ .

## 6 Applications and Related Results

The extended mean values and their generalizations have been applied not only to establish inequalities of the gamma function and the incomplete gamma function, to construct new Steffensen pairs, and to generalize the Hermite-Hadamard's inequality, but also to study quantum and to generalize the Bernoulli's numbers and polynomials.

### 6.1 Application to quantum

The concepts of the generalized weighted mean values  $M_{p,f}(r, s; x, y)$  have been applied to study of quantum in [49, 50].

## 6.2 Generalizations of Bernoulli's numbers and polynomials

The function  $g(t; x, y)$  defined by (1.12) has been applied to generalize the concepts of Bernoulli's numbers and polynomials. For details, please refer to [12, 22].

## 6.3 Generalization of Hermite-Hadamard's inequality

Using Tchebysheff's integral inequality, suitable properties of double integral and the revised Cauchy's mean value theorem in integral form of Lemma 5.1, the following result is proved.

**Theorem 6.1** ([13]). *Suppose  $f(x)$  is a positive differentiable function and  $w(x) \neq 0$  an integrable nonnegative weight on the interval  $[a, b]$ , if  $f'(x)$  and  $\frac{f'(x)}{w(x)}$  are integrable and both increasing or both decreasing, then, for all real numbers  $r$  and  $s$ , we have*

$$M_{w,f}(r, s; a, b) < E(r+1, s+1; f(a), f(b)); \quad (6.1)$$

if one of the functions  $f'(x)$  or  $\frac{f'(x)}{w(x)}$  is nondecreasing and the other nonincreasing, then inequality (6.1) reverses.

This inequality (6.1) generalizes Hermite-Hadamard's inequality. See [3, 13].

In [27], Hermite-Hadamard's inequality was generalized to the case of  $r$ -convex functions with the help of the extended mean values. In [21], the results obtained in [27] were further generalized to the case of so-called  $g$ -convex functions. Recently, it was further generalized and refined in [30].

## 6.4 Monotonicity results and inequalities involving gamma functions

It is well-known that the incomplete gamma function  $\Gamma(z, x)$  is defined for  $\text{Re } z > 0$  by (1.16) and define

$$\gamma(z, x) = \int_0^x t^{z-1} e^{-t} dt, \quad (6.2)$$

with  $\Gamma(z, 0) = \Gamma(z)$ , called the gamma function,  $\Gamma(0, x) = E_1(x)$  the exponential integral.

In [34], using inequality (6.1) and some results on the monotonicities of the generalized weighted mean values  $M_{p,f}(r, s; x, y)$ , it was verified that functions  $\left[\frac{\Gamma(s)}{\Gamma(r)}\right]^{1/(s-r)}$ ,  $\left[\frac{\Gamma(s, x)}{\Gamma(r, x)}\right]^{1/(s-r)}$  and  $\left[\frac{\gamma(s, x)}{\gamma(r, x)}\right]^{1/(s-r)}$  are increasing in  $r > 0$ ,  $s > 0$  and  $x > 0$ . From this, some

monotonicity results and inequalities for the gamma or the incomplete gamma functions are deduced or extended, a unified proof of some known results for the gamma function is given.

If taking  $p(t) = e^{-t}$  and  $f(t) = t$  for  $t \in (0, x)$  in Theorem 6.1, then we have

**Theorem 6.2 ([34]).** For fixed  $x > 0$ , the function  $\frac{s\gamma(s, x)}{x^s}$  is decreasing in  $s > 0$ .

From the monotonicity with the two parameters  $r$  and  $s$  of  $M_{p, f}(r, s; x, y)$  in Theorem 5.3, it follows that

**Theorem 6.3 ([34]).** The function  $\left[\frac{\Gamma(s)}{\Gamma(r)}\right]^{1/(s-r)}$  is increasing with  $r > 0$  and  $s > 0$ .

**Corollary 6.3.1 ([34]).** The functions  $[\Gamma(r)]^{1/(r-1)}$  and the digamma function  $\psi(r) = \frac{\Gamma'(r)}{\Gamma(r)}$ , the logarithmic derivative of the gamma function  $\Gamma(r)$ , are increasing in  $r > 0$ . Hence  $\Gamma(r)$  is a logarithmically convex function in the interval  $(0, \infty)$ .

*Remark 6.1.* In [18] and [23], among other things, the following monotonicity results were obtained

$$[\Gamma(1+k)]^{1/k} < [\Gamma(2+k)]^{1/(k-1)}, \quad k \in \mathbb{N};$$

$$\left[\Gamma\left(1 + \frac{1}{x}\right)\right]^x \text{ decreases with } x > 0.$$

Clearly, our Theorem 6.3 and Corollary 6.3.1 generalize and extend these results for the range of the argument.

**Corollary 6.3.2.** The following inequalities hold for  $s > r > 0$

$$\exp[(s-r)\psi(s)] > \frac{\Gamma(s)}{\Gamma(r)} > \exp[(s-r)\psi(r)], \quad (6.3)$$

$$e^{c\tau} < \Gamma(r+1) < \exp[r\psi(r+1)], \quad (6.4)$$

where  $c = 0.5772 \dots$  is the Euler's constant.

*Remark 6.2.* The ratio  $\frac{\Gamma(s)}{\Gamma(r)}$  has been researched by many mathematicians. W. Gautschi showed for  $0 < s < 1$  and  $n \in \mathbb{N}$  in [11] that

$$n^{1-s} < \frac{\Gamma(n+1)}{\Gamma(n+s)} < \exp[(1-s)\psi(n+1)]. \quad (6.5)$$

A strengthened upper bound was given by T. Erber in [7] as follows

$$\frac{\Gamma(n+1)}{\Gamma(n+s)} < \frac{4(n+s)(n+1)^{1-s}}{4n+(s+1)^2}, \quad 0 < s < 1, \quad n \in \mathbb{N}. \quad (6.6)$$

J. D. Kečkić and P. M. Vasić gave in [16] the inequalities below

$$\frac{b^{b-1}}{a^{a-1}} \cdot e^{a-b} < \frac{\Gamma(b)}{\Gamma(a)} < \frac{b^{b-1/2}}{a^{a-1/2}} \cdot e^{a-b}, \quad 0 < a < b. \quad (6.7)$$

The following closer bounds were proved for  $0 < s < 1$  and  $x \geq 1$  by D. Kershaw in [17].

$$\exp \left[ (1-s)\psi(x+s^{1/2}) \right] < \frac{\Gamma(x+1)}{\Gamma(x+s)} < \exp \left[ (1-s)\psi \left( x + \frac{s+1}{2} \right) \right], \quad (6.8)$$

$$\left( x + \frac{s}{2} \right)^{1-s} < \frac{\Gamma(x+1)}{\Gamma(x+s)} < \left[ x - \frac{1}{2} + \left( s - \frac{1}{4} \right)^{1/2} \right]^{1-s}. \quad (6.9)$$

It is easy to see that inequalities in (6.3) of Corollary 6.3.2 extend the range of arguments of above inequalities (6.5)–(6.9) but (6.7).

As consequences of Theorem 5.2 and Theorem 5.3, we have

**Theorem 6.4** ([34]). *For  $s > r > 0$  and  $x > 0$ , the functions  $\left[ \frac{\gamma(s,x)}{\gamma(r,x)} \right]^{1/(s-r)}$  and  $\left[ \frac{\Gamma(s,x)}{\Gamma(r,x)} \right]^{1/(s-r)}$  increase with either  $x$  or  $r$  and  $s$ . Therefore,  $\frac{\gamma(s,x)}{x^{s-1}}$  decreases and  $\frac{\Gamma(s,x)}{x^{s-1}}$  increases with  $s > 0$ , respectively.*

**Corollary 6.4.1.** *The incomplete gamma functions  $\gamma(r,x)$  and  $\Gamma(r,x)$  are logarithmically convex with respect to  $r > 0$  for fixed  $x > 0$ . The function  $\left[ \frac{\Gamma(r,x)}{E_1(x)} \right]^{1/r}$  is increasing in  $r > 0$  and  $x > 0$ . Therefore, the functions  $\frac{\Gamma(s+\theta)}{\Gamma(r+\theta)}$ ,  $\frac{\Gamma(s+\theta,x)}{\Gamma(r+\theta,x)}$  and  $\frac{\gamma(s+\theta,x)}{\gamma(r+\theta,x)}$  are increasing with  $\theta$  for fixed  $s > r > 0$  and  $x > 0$ .*

*Remark 6.3.* In the last week of November 2001, N. Elezović reminded me of his joint paper [5] with C. Giordana and J. Pečarić. In their paper [5], among others, the convexity with respect to variable  $x$  of the function  $\left[ \frac{\Gamma(x+t)}{\Gamma(x+s)} \right]^{1/(t-s)}$  for  $|t-s| < 1$  is verified, the best lower bound for (6.8) and the best upper bound for (6.9) are obtained, some different approach from Gautschi's in [11] is given, several new simple inequalities for digamma function are also proved.

The gamma and incomplete gamma functions and related functions have been investigated using different approaches, for examples, see [1, 4, 38, 40, 41, 43].

## 6.5 Establishment of Steffensen pairs

Let  $f$  and  $g$  be integrable functions on  $[a, b]$  such that  $f$  is decreasing and  $0 \leq g(x) \leq 1$  for  $x \in [a, b]$ . Then

$$\int_{b-\lambda}^b f(x) dx \leq \int_a^b f(x)g(x) dx \leq \int_a^{a+\lambda} f(x) dx, \quad (6.10)$$

where  $\lambda = \int_a^b g(x) dx$ .

The inequality (6.10) is called Steffensen's inequality.

In [8], a discrete analogue of the inequality (6.10) was proved: Let  $\{x_i\}_{i=1}^n$  be a decreasing finite sequence of nonnegative real numbers,  $\{y_i\}_{i=1}^n$  be a finite sequence of real numbers such that  $0 \leq y_i \leq 1$  for  $1 \leq i \leq n$ . Let  $k_1, k_2 \in \{1, 2, \dots, n\}$  be such that  $k_2 \leq \sum_{i=1}^n y_i \leq k_1$ . Then

$$\sum_{i=n-k_2+1}^n x_i \leq \sum_{i=1}^n x_i y_i \leq \sum_{i=1}^{k_1} x_i. \quad (6.11)$$

As a direct consequence of inequality (6.11), we have: Let  $\{x_i\}_{i=1}^n$  be nonnegative real numbers such that  $\sum_{i=1}^n x_i \leq A$  and  $\sum_{i=1}^n x_i^2 \geq B^2$ , where  $A$  and  $B$  are positive real numbers. Let  $k \in \{1, 2, \dots, n\}$  be such that  $k \geq \frac{A}{B}$ . Then there are  $k$  numbers among  $x_1, x_2, \dots, x_n$  whose sum is bigger than or equals to  $B$ .

The so-called Steffensen pair was defined by H. Gauchman in [10] as follows.

**Definition 6.1.** Let  $\varphi: [c, \infty) \rightarrow [0, \infty)$  and  $\tau: (0, \infty) \rightarrow (0, \infty)$  be two strictly increasing functions,  $c \geq 0$ , let  $\{x_i\}_{i=1}^n$  be a finite sequence of real numbers such that  $x_i \geq c$  for  $1 \leq i \leq n$ ,  $A$  and  $B$  be positive real numbers, and  $\sum_{i=1}^n x_i \leq A$ ,  $\sum_{i=1}^n \varphi(x_i) \geq \varphi(B)$ . If, for any  $k \in \{1, 2, \dots, n\}$  such that  $k \geq \tau(\frac{A}{B})$ , there are  $k$  numbers among  $x_1, \dots, x_n$  whose sum is not less than  $B$ , then we call  $(\varphi, \tau)$  a Steffensen pair on  $[c, \infty)$ .

The following Steffensen pairs were found by H. Gauchman in [10].

$$(x^\alpha, x^{1/(\alpha-1)}), \quad \alpha \geq 2, \quad x \in [0, \infty); \quad (6.12)$$

$$(x \exp(x^\alpha - 1), (1 + \ln x)^{1/\alpha}), \quad \alpha \geq 1, \quad x \in [1, \infty). \quad (6.13)$$

Let  $a$  and  $b$  be real numbers satisfying  $b > a > 1$  and  $\sqrt{ab} \geq e$ . Define

$$\varphi(x) = \begin{cases} \frac{x^{1+\ln b} - x^{1+\ln a}}{\ln x} & \text{if } x > 1, \\ \ln b - \ln a & \text{if } x = 1, \end{cases} \quad (6.14)$$

$$\tau(x) = x^{1/\ln \sqrt{ab}}. \quad (6.15)$$

Then it was verified by H. Gauchman in [10] that  $(\varphi, \tau)$  is a Steffensen pair on  $[1, \infty)$  using some results and techniques in [46].

With help of properties of the extended mean values  $E(r, s; x, y)$  and the generalized weighted mean values  $M_{p,f}(r, s; x, y)$ , some new Steffensen pairs were established in [37, 39].

Using the integral expression (1.14) of function  $\frac{b^x - a^x}{x}$ , mathematical induction and analytic techniques, we have

**Theorem 6.5 ([37]).** *If  $a$  and  $b$  are real numbers satisfying  $b > a > 1$  or  $b > \frac{1}{a} > 1$ , and  $\sqrt{ab} \geq e$ , then*

$$\left( x \int_a^b t^{\ln x - 1} dt, x^{2/\ln(ab)} \right) \quad (6.16)$$

is a Steffensen pair on  $[1, \infty)$ . If  $a$  and  $b$  are real numbers satisfying  $b > a > 1$  and  $\sqrt{ab} \geq e$ , then

$$\left( x \int_a^b (\ln t)^n t^{\ln x - 1} dt, x^{\frac{n+2}{n-1} \frac{(\ln b)^{n+1} - (\ln a)^{n+1}}{(\ln b)^{n+2} - (\ln a)^{n+2}}} \right) \quad (6.17)$$

are Steffensen pairs on  $[1, \infty)$  for any positive integer  $n$ .

In [39], considering the function  $\int_a^b p(u) f^t(u) du$  and its properties, we further obtain more general Steffensen pairs as follows.

**Theorem 6.6 ([39]).** *Let  $a, b \in \mathbb{R}$ , let  $p \neq 0$  be a nonnegative and integrable function and  $f$  a positive and integrable function on the interval  $[a, b]$ .*

1. *If inequality*

$$\int_a^b p(u) du \leq \int_a^b p(u) \ln f(u) du \quad (6.18)$$

holds, then

$$\left( x \int_a^b p(u) [f(u)]^{\ln x} du, x^{\frac{\int_a^b p(u) du}{\int_a^b p(u) \ln f(u) du}} \right) \quad (6.19)$$

is a Steffensen pair on  $[1, \infty)$ .

2. *If  $f(u) \geq 1$  and inequality (6.18) holds, then*

$$\left( x \int_a^b p(u) [f(u)]^{\ln x} [\ln f(u)]^n du, x^{\frac{\int_a^b p(u) \ln f(u) du}{\int_a^b p(u) [\ln f(u)]^{n+1} du}} \right) \quad (6.20)$$

are Steffensen pairs on  $[1, \infty)$  for any positive integer  $n$ .



## Acknowledgements

This paper was completed during the author's visit to the RGMIA between November 1, 2001 and January 31, 2002, as a Visiting Professor with grants from the Victoria University of Technology and Jiaozuo Institute of Technology.

## References

- [1] ALLASIA, G., GIORDANO, C. AND PEČARIĆ, J., *Hadamard-type inequalities for  $(2r)$ -convex functions with applications*, Atti Accad. Sci. Torino Cl. Sci. Fis Mat Natur. **133** (1999), 187–200.
- [2] BECKENBACH, E.F. AND BELLMAN, R., *Inequalities*. Springer, Berlin, 1983.
- [3] DRAGOMIR, S.S. AND PEARCE, C.E.M., *Selected Topics on Hermite-Hadamard Type Inequalities and Applications*, RGMIA Monographs. (2000) Available online at [http://rgmia.vu.edu.au/monographs/hermite\\_hadamard.html](http://rgmia.vu.edu.au/monographs/hermite_hadamard.html)
- [4] ELBERT, Á. AND LAFORGIA, A., *An inequality for the product of two integrals relating to the incomplete gamma function*, J. Inequal. Appl. **5** (2000), 39–51.
- [5] ELEZOVIĆ, N., GIORDANO, C. AND PEČARIĆ, J., *The best bounds in Gautschi's inequality*, Math. Inequal. Appl. **3** (2000), no. 2, 239–252.
- [6] ELEZOVIĆ, N. AND PEČARIĆ, J., *A note on Schur-convex functions*, Rocky Mountain J. Math. **30** (2000), no. 3, 853–856.
- [7] ERBER, T., *The gamma function inequalities of Gurland and Gautschi*, Scand. Actuar. J. **1960** (1961), 27–28.
- [8] EVARD, J.-C. AND GAUCHMAN, H., *Steffensen type inequalities over general measure spaces*, Analysis **17** (1997), 301–322.
- [9] FINK, A.M., *An essay on the history of inequalities*, J. Math. Anal. Appl. **249** (2000), 118–134.
- [10] GAUCHMAN, H., *Steffensen pairs and associated inequalities*, J. Inequal. Appl. **5** (2000), no. 1, 53–61.
- [11] GAUTSCHI, W., *Some elementary inequalities relating to the gamma and incomplete gamma function*, J. Math. Phys. **38** (1959), 77–81.
- [12] GUO, B.-N. AND QI, F., *Generalisation of Bernoulli polynomials*, Internat. J. Math. Ed. Sci. Tech. (2002), in the press. RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 10, 691–695. Available online at <http://rgmia.vu.edu.au/v4n4.html>.

- [13] GUO, B.-N. AND QI, F., *Inequalities for generalized weighted mean values of convex function*, Math. Inequal. Appl. **4** (2001), no. 2, 195–202.
- [14] GUO, B.-N. AND QI, F., *Proofs of an integral inequality*, Mathematics and Informatics Quarterly **7** (1997), no. 4, 182–184.
- [15] GUO, B.-N., ZHANG, SH.-Q. AND QI, F., *Elementary proofs of monotonicity for extended mean values of some functions with two parameters*, Shùxué de Shíjiān yǔ Rènshì (Mathematics in Practice and Theory) **29** (1999), no.2, 169–174, (Chinese).
- [16] KEČKIĆ, J.D. AND VASIĆ, P.M., *Some inequalities for the gamma function*, Publ. Inst. Math. Beograd N. S. **11** (1971), 107–114.
- [17] KERSHAW, D., *Some extensions of W. Gautschi's inequalities for the gamma function*, Math. Comp. **41** (1983), 607–611.
- [18] KERSHAW, D. AND LAFORGIA, A., *Monotonicity results for the gamma function*, Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **119** (1985), 127–133.
- [19] LEACH, E.B. AND SHOLANDER, M.C., *Extended mean values*, Amer. Math. Monthly **85** (1978), 84–90.
- [20] LEACH, E.B. AND SHOLANDER, M.C., *Extended mean values II*, J. Math. Anal. Appl. **92** (1983), 207–223.
- [21] LEE, K.-CH. AND TSENG, K.-L., *On weighted generalization of Hadamard's inequality for  $g$ -convex functions*, Tamsui Oxf. J. Math. Sci. **16** (2000), no. 1, 91–104.
- [22] LUO, Q.-M., GUO, B.-N. AND QI, F., *Generalizations of Bernoulli numbers and polynomials*, submitted.
- [23] MINC, H. AND SATHRE, L., *Some inequalities involving  $(r!)^{1/r}$* , Proc. Edinburgh Math. Soc. **14** (1965/66), 41–46.
- [24] MITRINOVIĆ, D.S., *Analytic Inequalities*, Springer-Verlag, New York/Heidelberg/Berlin, 1970.
- [25] MITRINOVIĆ, D.S., PEČARIĆ, J.E. AND FINK, A.M., *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.
- [26] PÁLES, Z., *Inequalities for differences of powers*, J. Math. Anal. Appl. **131** (1988), 271–281.
- [27] PEARCE, C.E.M., PEČARIĆ, J. AND ŠIMIĆ, V., *Stolarsky means and Hadamard's inequality*, J. Math. Anal. Appl. **220** (1998), 99–109.
- [28] PEČARIĆ, J., PROSCHAN, F. AND TONG, Y.L., *Convex Functions, Partial Orderings, and Statistical Applications*, Mathematics in Science and Engineering **187**, Academic Press, 1992.

- [29] PEČARIĆ, J., QI, F., ŠIMIĆ, V. AND XU, S.-L., *Refinements and extensions of an inequality, III*, J. Math. Anal. Appl. **227** (1998), no. 2, 439–448.
- [30] QI, F., *Generalizations and refinements of Hermite-Hadamard's inequality*, submitted.
- [31] QI, F., *Generalized abstracted mean values*, J. Inequal. Pure Appl. Math. **1** (2000), no. 1, Art. 4. Available online at [http://jipam.vu.edu.au/v1n1/013\\_99.html](http://jipam.vu.edu.au/v1n1/013_99.html). RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 4, 633–642. Available online at <http://rgmia.vu.edu.au/v2n5.html>.
- [32] QI, F., *Generalized weighted mean values with two parameters*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **454** (1998), no. 1978, 2723–2732.
- [33] QI, F., *Logarithmic convexity of extended mean values*, Proc. Amer. Math. Soc. **130** (2002), no. 6, 1787–1796. <http://www.ams.org/journal-getitem?pii=S0002-9939-01-06275-X>. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 5, 643–652. Available online at <http://rgmia.vu.edu.au/v2n5.html>.
- [34] QI, F., *Monotonicity results and inequalities for the gamma and incomplete gamma functions*, Math. Inequal. Appl. **5** (2002), no. 1, 61–67. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 7. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [35] QI, F., *On a two-parameter family of nonhomogeneous mean values*, Tamkang J. Math. **29** (1998), no. 2, 155–163.
- [36] QI, F., *Schur-convexity of the extended mean values*, RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 4. Available online at <http://rgmia.vu.edu.au/v4n4.html>.
- [37] QI, F., CHENG, J.-X. AND WANG, G., *New Steffensen pairs*, Proceedings of the 6th International Conference 2000 on Nonlinear Functional Analysis and Applications: Inequality Theory and Applications **1** (2002), 279–285. RGMIA Res. Rep. Coll. **3** (2000), no. 3, Art. 11. Available online at <http://rgmia.vu.edu.au/v3n3.html>.
- [38] QI, F., CUI, L.-H. AND XU, S.-L., *Some inequalities constructed by Tchebysheff's integral inequality*, Math. Inequal. Appl. **2** (1999), no. 4, 517–528.
- [39] QI, F. AND GUO, B.-N., *On Steffensen pairs*, J. Math. Anal. Appl. (2002), in the press. RGMIA Res. Rep. Coll. **3** (2000), no. 3, Art. 10, 425–430. Available online at <http://rgmia.vu.edu.au/v3n3.html>.
- [40] QI, F. AND GUO, B.-N., *Some inequalities involving the geometric mean of natural numbers and the ratio of gamma functions*, RGMIA Res. Rep. Coll. **4** (2001), no. 1, Art. 6, 41–48. Available online at <http://rgmia.vu.edu.au/v4n1.html>.
- [41] QI, F. AND GUO, S.-L., *Inequalities for the incomplete gamma and related functions*, Math. Inequal. Appl. **2** (1999), no. 1, 47–53.

- [42] QI, F. AND LUO, Q.-M., *A simple proof of monotonicity for extended mean values*, J. Math. Anal. Appl. **224** (1998), no. 2, 356–359.
- [43] QI, F. AND MEI, J.-Q., *Some inequalities for the incomplete gamma and related functions*, Z. Anal. Anwendungen **18** (1999), no. 3, 793–799.
- [44] QI, F., MEI, J.-Q., XIA, D.-F. AND XU, S.-L., *New proofs of weighted power mean inequalities and monotonicity for generalized weighted mean values*, Math. Inequal. Appl. **3** (2000), no. 3, 377–383.
- [45] QI, F., SÁNDOR, J., DRAGOMIR, S.S. AND SOFO, A., *Notes on the Schur-convexity of the extended mean values*, RGMIA Res. Rep. Coll. **5** (2002), no. 1, Art. 3. Available online at <http://rgmia.vu.edu.au/v5n1.html>.
- [46] QI, F. AND XU, S.-L., *The function  $(b^x - a^x)/x$ : Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359. <http://www.ams.org/journal-getitem?pii=S0002-9939-98-04442-6>.
- [47] QI, F., XU, S.-L. AND DEBNATH, L., *A new proof of monotonicity for extended mean values*, Internat. J. Math. Math. Sci. **22** (1999), no. 2, 415–420.
- [48] QI, F. AND ZHANG, SH.-Q., *Note on monotonicity of generalized weighted mean values*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **455** (1999), no. 1989, 3259–3260.
- [49] SLATER, P.B., *A priori probabilities of separable quantum states*, J. Phys. A-Math. Gen. **32** (1999), no. 28, 5261–5275.
- [50] SLATER, P.B., *Hall normalization constants for the Bures volumes of the  $n$ -state quantum systems*, J. Phys. A-Math. Gen. **32** (1999), no. 47, 8231–8246.
- [51] STOLARSKY, K.B., *Generalizations of the logarithmic mean*, Mag. Math. **48** (1975), 87–92.
- [52] WIDDER, D.V., *The Laplace Transform*, Princeton University Press, Princeton, 1941.